

M.C. González-García · Michele Maltoni · Joan Rojo

Determination of the atmospheric neutrino fluxes from experimental data

Received: date / Accepted: date

Abstract The precise knowledge of the atmospheric neutrino fluxes is a key ingredient in the interpretation of the results from any atmospheric neutrino experiment. In the standard atmospheric neutrino data analysis, these fluxes are theoretical inputs obtained from sophisticated numerical calculations. In this work we present an alternative approach to the determination of the atmospheric neutrino fluxes based on the direct extraction from the experimental data on neutrino event rates.

Keywords atmospheric neutrinos · neutrino oscillations

PACS 14.60.Lm · 14.60.Pq

1 Introduction

One of the most important breakthroughs in particle physics, and the only solid evidence for physics beyond the Standard Model, is the discovery – following a variety of independent experiments [11] – that neutrinos are massive and consequently can oscillate among their different flavor eigenstates. The flavor oscillation hypothesis has been supported by an impressive wealth of experimental data, one of the most important pieces of evidence coming from atmospheric neutrinos.

M. C. González-García
C.N. Yang Institute for Theoretical Physics
State University of New York at Stony Brook
Stony Brook, NY 11794-3840, USA,
and IFIC, Universitat de València – C.S.I.C.
Edificio Institutos de Paterna, Apt 22085, 46071 València,
Spain
E-mail: concha@insti.physics.sunysb.edu

M. Maltoni
International Centre for Theoretical Physics,
Strada Costiera 11, 31014 Trieste, Italy
E-mail: mmaltoni@ictp.trieste.it

J. Rojo
Departament d'Estructura i Constituents de la Matèria
Diagonal 647, 08028 Barcelona, Spain
E-mail: joanrojo@ecm.ub.es

Atmospheric neutrinos are originated by the collisions of cosmic rays with air nuclei in the Earth atmosphere. In the collision mostly pions (and some kaons) are produced and they subsequently decay into electron and muon neutrinos and anti-neutrinos. These neutrinos are observed in underground experiments using different techniques [1]. In particular in the last ten years, high precision and large statistics data has been available from the SuperKamiokande experiment [1] which has clearly established the existence of a deficit in the μ -like atmospheric events with the expected distance and energy dependence from $\nu_\mu \rightarrow \nu_\tau$ oscillations with oscillation parameters $\Delta m^2 \sim 2 \times 10^{-3} \text{ eV}^2$ and $\tan^2 \theta = 1$. This evidence has also been confirmed by other atmospheric experiments such as MACRO and Soudan 2.

The expected number of atmospheric neutrino events depends on a variety of components: the atmospheric neutrino fluxes, the neutrino oscillation parameters and the neutrino-nucleus interaction cross section. Since the main focus of atmospheric neutrino data interpretation has been the determination of neutrino oscillation parameters, in the standard analysis the remaining components of the event rate computation are inputs taken from other sources. In particular, the fluxes of atmospheric neutrinos are taken from the results of numerical calculations, like those of Refs. [13, 2], which make use of a mixture of primary cosmic ray spectrum measurements, models for the hadronic interactions and simulation of particle propagation [7].

The attainable accuracy in the independent determination of the relevant neutrino oscillation parameters from non-atmospheric neutrino experiments makes it possible to attempt an inversion of the strategy in the atmospheric neutrino analysis: to use the oscillation parameters as inputs in the data analysis in order to extract from data the atmospheric neutrino fluxes.

There are several motivations for such direct determination of the atmospheric neutrino flux from experimental data. First of all it would provide a cross-check of the standard flux calculations as well as of the size of the associated uncertainties (which being mostly theoret-

ical are difficult to quantify). Second, a precise knowledge of atmospheric neutrino flux is of paramount importance for high energy neutrino telescopes [12], both because they are the main background and they are used for detector calibration. Finally, such program will quantitatively expand the physics potential of future atmospheric neutrino experiments. Technically, however, this program is challenged by the absence of a generic parametrization of the energy and angular functional dependence of the fluxes which is valid in all the range of energies where there is available neutrino data.

In this contribution we present the first results on this alternative approach to the determination of the atmospheric neutrino fluxes: we will determine these fluxes from experimental data on atmospheric neutrino event rates, using all available information on neutrino oscillation parameters, cross-sections, as well as information from the different sources of experimental and theoretical uncertainty. The problem of the unknown functional form for the neutrino flux is bypassed by the use of neural networks as interpolants, since they allow us to parametrize the atmospheric neutrino flux without having to assume any functional behavior. Additional details on our approach can be found in [9].

Indeed the problem of the *deconvolution* of the atmospheric flux from experimental data on event rates is rather close in spirit to the determination of parton distribution functions in deep-inelastic scattering from experimentally measured structure functions [5]. For this reason, in this work we will apply for the determination of the atmospheric neutrino fluxes a general strategy originally designed to extract parton distributions in an unbiased way with faithful estimation of the uncertainties¹ [6,3,4].

2 General strategy

The general strategy that will be used to determine the atmospheric neutrino fluxes was first presented in Ref. [6] (see also [15]). It involves two distinct stages in order to go from the data to the flux parametrization. In the first stage, a Monte Carlo sample of replicas of the experimental data on neutrino event rates (“artificial data”) is generated. These can be viewed as a sampling of the probability measure on the space of physical observables at the discrete points where data exist. In the second stage one uses neural networks to interpolate between points where data exist. In the present case, this second stage in turn consists of two sub-steps: the determination of the atmospheric event rates from the atmospheric flux in a fast and efficient way, and the comparison of the event rates thus computed to the data in order to tune the best-fit form of input neural flux distribution

¹ This strategy has also been successfully applied with different motivations in other contexts like tau lepton decays [17] and B meson physics [16]

(“training of the neural network”). Combining these two steps, the space of physical observables is mapped onto the space of fluxes, so the experimental information on the former can be interpolated by neural networks in the latter.

In the present analysis we use the latest data on atmospheric neutrino event rates from the Super Kamiokande Collaboration [1]. Higher energy data from neutrino telescopes like Amanda [8] is not publicly available in a format which allows for its inclusion in the present analysis and its treatment is left for future work.

The latest Super Kamiokande atmospheric neutrino data sample is divided in 9 different types of events: contained events in three energy ranges, Sub-GeV, Mid-GeV and Multi-GeV electron- and muon-like, partially contained muon-like events and upgoing stopping and throughgoing muon events. Each of the above types of events is divided in 10 bins in the final state lepton zenith angle ϕ_l . Therefore we have a total of $N_{\text{dat}} = 90$ experimental data points, which we label as

$$R_i^{(\text{exp})}, \quad i = 1, \dots, N_{\text{dat}}. \quad (1)$$

Note that each type of atmospheric neutrino event rate is sensitive to a different region of the neutrino energy spectrum. In particular we note that for energies larger than $E_\nu \sim \text{few TeV}$ and smaller than $E_\nu \sim 0.1 \text{ GeV}$ there is essentially no information on the atmospheric flux from the available event rates.

The purpose of the artificial data generation is to produce a Monte Carlo set of ‘pseudo-data’, i.e. N_{rep} replicas of the original set of N_{dat} data points, $R_i^{(\text{art}) (k)}$ such that the N_{rep} sets of N_{dat} points are distributed according to an N_{dat} -dimensional multi-gaussian distribution around the original points, with expectation values equal to the central experimental values, and error and covariance equal to the corresponding experimental quantities. This is achieved by defining

$$R_i^{(\text{art}) (k)} = R_i^{(\text{exp})} + r_i^{(k)} \sigma_i^{\text{tot}}, \quad k = 1, N_{\text{rep}}, \quad (2)$$

where N_{rep} is the number of generated replicas of the experimental data, and where $r_i^{(k)}$ are univariate gaussian random numbers with the same correlation matrix as experimental data. We can then generate arbitrarily many sets of pseudo-data, and choose the number of sets N_{rep} in such a way that the properties of the Monte Carlo sample reproduce those of the original data set to arbitrary accuracy.

In our case each neural network parametrizes a neutrino flux. Since the precision of the available experimental data is not enough to allow for a separate determination of the energy, zenith angle and type dependence of the atmospheric flux, in this work we will assume the zenith and type dependence of the flux to be known with some precision and extract from the data only its energy dependence. That is, if $\text{NN}(E_\nu)$ is the neural network

output when the input is the neutrino energy E_ν , then the neural flux parametrization will be

$$\Phi^{(\text{net})}(E_\nu, c_\nu, t) = \text{NN}(E_\nu) \Phi^{(\text{ref})}(E_\nu, c_\nu, t) . \quad (3)$$

$\Phi^{(\text{ref})}$ is a reference differential flux, which we take to be the most recent computations of either the Honda [13] or the Bartol [2] collaborations, which have been extended to cover also the high-energy region by consistent matching with the Volkova fluxes.

Now let us describe a fast and efficient technique to evaluate the atmospheric neutrino event rates $R_i^{(\text{net})}$ for an arbitrary input neural network atmospheric flux $\Phi^{(\text{net})}(E_\nu, c_\nu, t)$. For example, for contained events any rate $R_i^{(\text{net})}$ can be computed

$$R_i^{(\text{net})} = n_t T \sum_{\alpha, \beta, \pm} \int_0^\infty dh \int_{-1}^{+1} dc_\nu \int_{E_{\min}}^\infty dE_\nu \int_{E_{\min}}^{E_\nu} dE_l \int_{-1}^{+1} dc_a \int_0^{2\pi} d\varphi_a \frac{d^3 \Phi_\alpha^{\pm(\text{net})}}{dE_\nu dc_\nu dh}(E_\nu, c_\nu, h) \\ P_{\alpha \rightarrow \beta}^\pm(E_\nu, c_\nu, h | \boldsymbol{\eta}) \frac{d^2 \sigma_\beta^\pm}{dE_l dc_a}(E_\nu, E_l, c_a) \varepsilon_\beta^{\text{bin}}(E_l, c_l) , \quad (4)$$

where Φ_α^+ (Φ_α^-) is the flux of atmospheric neutrinos (antineutrinos) of type α , σ_β^+ (σ_β^-) is the charged-current neutrino- (antineutrino-) nucleon interaction cross section, and see [10] for a more detailed description of the above expression. The above expressions are very time-consuming to be evaluated, and this is specially a problem in our case since in the neural network approach one requires a very large number of evaluations.

However, the procedure can be speed up if one realizes that for a given flux, the expected event rates can always be written as

$$R_i = \sum_t \int_{-1}^1 dc_\nu \int_{E_{\min}}^\infty dE_\nu \Phi(E_\nu, c_\nu, t) \tilde{C}_i(E_\nu, c_\nu, t) , \quad (5)$$

where $\tilde{C}_i(E_\nu, c_\nu, t)$ contain all the E_ν and c_ν flux independent pieces in Eq.(4). Consequently if one discretizes the differential fluxes in N_e energy intervals and N_z zenith angle intervals

$$\Phi^{(\text{net})}(E_\nu, c_\nu, t) = \sum_{e,z} \Psi_{e z t}^{(\text{net})} \theta(E_\nu \in I_e) \theta(c_\nu \in I_z) , \quad (6)$$

it is possible to write the theoretical predictions as a sum of the elements of a bin-integrated flux table $\Psi_{e z t}^{(\text{net})}$,

$$R_i^{(\text{net})} = \sum_{e z t} C_{e z t}^i \Psi_{e z t}^{(\text{net})} , \quad (7)$$

where the coefficients $C_{e z t}^i$, which are the most time-consuming ingredient, need only to be precomputed once

before the training, since they do not depend on the parametrization of the atmospheric neutrino flux.

The determination of the parameters that define the neural network, its weights, is performed by maximum likelihood. This procedure, the so-called neural network training, proceeds by minimizing an error function, which coincides with the χ^2 of the experimental points when compared to their theoretical determination obtained using the given set of fluxes:

$$E^{(k)}(\{\omega_i\}) = \min_{\boldsymbol{\xi}} \left[\sum_{i, \text{theo}} \xi_i^2 + \sum_{i, \text{sys}} \xi_i^2 \right] \quad (8)$$

$$\left(\sum_{n=1}^{N_{\text{dat}}} \frac{R_n^{(\text{net})(k)}(\{\omega_i\}) \left[1 + \sum_i \pi_i^n \xi_i \right] - R_n^{(\text{art})(k)}}{\sigma_n^{\text{stat}}} \right)^2 \quad (9)$$

The error function has to be minimized with respect to $\{\omega_i\}$, the parameters of the neural network. The minimization of Eq. (8) is performed with the use of genetic algorithms [15]. Because of the nonlinear dependence of the neural net on its parameters, and the nonlocal dependence of the measured quantities on the neural net (event rates are given by multi-dimensional convolutions of the initial flux distributions), a genetic algorithm turns out to be the most efficient minimization method.

Thus at the end of the procedure, we end up with N_{rep} fluxes, with each flux $\Phi^{(\text{net})(k)}$ given by a neural net. The set of N_{rep} fluxes provide our best representation of the corresponding probability density in the space of atmospheric neutrino fluxes: for example, the mean value of the flux at a given value of E_ν is found by averaging over the replicas, and the uncertainty on this value is the variance of the values given by the replicas.

3 Results and discussion

Now we discuss the results of our determination of the atmospheric neutrino fluxes. Note that any functional of the neutrino flux (like for example event rates and its associated uncertainty) can be computed taking the appropriate moments over the constructed probability measure

$$\langle \mathcal{F}(\Phi(E_\nu)) \rangle_{\text{rep}} = \frac{1}{N_{\text{rep}}} \sum_{k=1}^{N_{\text{rep}}} \mathcal{F}(\Phi^{(\text{net})(k)}(E_\nu)) . \quad (10)$$

In Fig. 1 we show our results for the atmospheric neutrino flux as compared with the Honda and Bartol computations as well as with some direct measurements from Amanda [8]. In Fig. 2 we show that the effects of the uncertainties in the neutrino oscillation parameters are




Fig. 1 Comparison of the $1 - \sigma$ uncertainty band of our parametrization of the atmospheric neutrino flux with recent computations and AMANDA data.




Fig. 2 Results for fits varying the default neutrino oscillation parameters from their central values within their $1 - \sigma$ uncertainties [14].

rather small with respect the intrinsic flux uncertainties. Finally, we show in Fig. 3 how our neural network parametrization successfully reproduces the features of experimental data.

The work presented in this contribution can be extended in several directions. First of all one could reduce the uncertainty in the large E_ν region by incorporating in the fit atmospheric neutrino measurements from neutrino detectors like Amada [8]. Second, we could use at even larger energies simple parametrizations of the flux to obtain a better estimation of backgrounds at neutrino telescopes. Finally, we could estimate the required statistics that would be required in forthcoming atmospheric neutrino experiments in order to determine from data also the zenith angle and flavor dependence of the neutrino flux.




Fig. 3 Comparison of the neural network results with experimental data for the Multi-GeV ν_μ data sample.

References

1. Ashie, Y., et al.: A measurement of atmospheric neutrino oscillation parameters by super-kamiokande i. *Phys. Rev. D* **71**, 112,005 (2005)
2. Barr, G.D., Gaisser, T.K., Lipari, P., Robbins, S., Stanev, T.: A three-dimensional calculation of atmospheric neutrinos. *Phys. Rev. D* **70**, 023,006 (2004)
3. Del Debbio, L., Forte, S., Latorre, J.I., Piccione, A., Rojo, J.: Unbiased determination of the proton structure function $f_2(p)$ with faithful uncertainty estimation. *JHEP* **03**, 080 (2005)
4. Del Debbio, L., Forte, S., Latorre, J.I., Piccione, A., Rojo, J.: The neural network approach to parton distribution functions: The nonsinglet case (In preparation)
5. Ellis, R.K., Stirling, W.J., Webber, B.R.: Qcd and collider physics. *Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol.* **8**, 1–435 (1996)
6. Forte, S., Garrido, L., Latorre, J.I., Piccione, A.: Neural network parametrization of deep-inelastic structure functions. *JHEP* **05**, 062 (2002)
7. Gaisser, T.K., Honda, M.: Flux of atmospheric neutrinos. *Ann. Rev. Nucl. Part. Sci.* **52**, 153–199 (2002)
8. Geenen, H.: Atmospheric neutrino and muon spectra measured with the amanda-ii detector (2003). Prepared for 28th International Cosmic Ray Conferences (ICRC 2003), Tsukuba, Japan, 31 Jul - 7 Aug 2003
9. Gonzalez-Garcia, C., Maltoni, M., Rojo, J.: Determination of atmospheric neutrino fluxes from atmospheric neutrino data. *arxiv: hep-ph/0607324*.
10. Gonzalez-Garcia, M.C., Maltoni, M.: Atmospheric neutrino oscillations and new physics. *Phys. Rev. D* **70**, 033,010 (2004)
11. Gonzalez-Garcia, M.C., Nir, Y.: Developments in neutrino physics. *Rev. Mod. Phys.* **75**, 345–402 (2003)
12. Halzen, F.: Lectures on high-energy neutrino astronomy (2005)
13. Honda, M., Kajita, T., Kasahara, K., Midorikawa, S.: A new calculation of the atmospheric neutrino flux in a 3-dimensional scheme. *Phys. Rev. D* **70**, 043,008 (2004)
14. Maltoni, M., Schwetz, T., Tortola, M.A., Valle, J.W.F.: Status of global fits to neutrino oscillations. *New J. Phys.* **6**, 122 (2004)
15. Rojo, J.: The neural network approach to parton distribution functions. Ph.D. thesis, Universitat de Barcelona (2006)
16. Rojo, J.: Neural network parametrization of the lepton energy spectrum in semileptonic b meson decays. *JHEP* **05**, 040 (2006)

-
17. Rojo, J., Latorre, J.I.: Neural network parametrization of spectral functions from hadronic tau decays and determination of qcd vacuum condensates. *JHEP* **01**, 055 (2004)